

DIVERGENT SELECTION FOR INTRAMUSCULAR FAT IN RABBITS. DIRECT AND CORRELATED RESPONSES TO SELECTION. PRELIMINARY RESULTS.

Blasco A., Zomeño C., Hernández P.

Institute for Animal Science and Technology, Universitat Politècnica de València, Camino de Vera s/n. P.O.Box 22012, 46022, Valencia, Spain.

*Corresponding author: crizose@posgrado.upv.es

ABSTRACT

The objective of this study was to evaluate the genetic response for a divergent selection experiment on intramuscular fat (IMF) content and to determine the correlated response in carcass and meat quality traits. Data from two generations of selection were used in this study. The base population was composed by 13 males and 83 females. High (H) and Low (L) lines had 8 males and approximately 40 females each one per generation. Selection was based on the average phenotypic IMF value of two full sibs measured in *Longissimus* muscle (LM). Two rabbits (male and female) of the first parity of each doe were slaughtered at 9 weeks of age. The following traits were recorded: live weight (LW), hot carcass weight, chilled carcass weight, reference carcass weight, scapular fat weight, perirenal fat weight, meat/bone ratio of hind leg, pH of LM, colour (lightness, L*; redness, a*; yellowness, b*) of the carcass and LM. *Longissimus* muscles were excised from the carcass and the IMF content was measured by near infrared reflectance spectroscopy. Data were analyzed using Bayesian methodology. Response to selection was estimated by the difference between H and L lines. The difference between lines for IMF content in the second generation of selection was 0.11 g/100g, which represents a direct response to selection of 9.8 %, of which a 6.8 % was obtained in the first generation and a 3 % in the second. It seems to be a negative correlated response on live weight, carcass and adipose tissues weights, with higher values in L line. However, the evidence for adipose tissue is rather weak. There is no evidence of changes in carcass and meat colour measurements as well as in meat/bone ratio and muscle pH.

Key words: intramuscular fat, near infrared reflectance spectroscopy, response, selection.

INTRODUCTION

Intramuscular fat (IMF) content is generally considered one of the main factors affecting meat quality (Wood et al., 2008). IMF can be improved by selection due to its high heritability (an average of 0.5 in pigs; Sellier 1998 and Suzuki et al., 2005a) and variability. However, there is scarce information about direct and correlated responses to selection for IMF, only three studies tried to increase IMF by selection in cattle (Sapp et al., 2002) and pigs (Suzuki et al. 2005b; Schwab et al., 2009).

Genetic programmes need a substantial amount of data to obtain estimates of genetic parameters with a minimal precision, thus genetic correlations involving meat quality analysis are often not accurate. Divergent selection experiments can be an alternative to examine the genetic

determination of the traits of interest. Direct and correlated responses can be studied by comparing the means of high and low lines for different traits after selection, which requires fewer individuals to obtain reasonable accuracies. Rabbit is an excellent model for genetic studies in other species due to its reduced generation interval and the low cost of the carcasses.

The objective of this study was to evaluate the direct response to selection on IMF content and to determine the correlated response in carcass and meat quality traits in rabbits.

MATERIALS AND METHODS

Animals and experimental design

A total of 502 rabbits from a divergent selection experiment for intramuscular fat content (IMF) were used in this study. Animals came from a synthetic line selected for ovulation rate (Laborda et al., 2011), and then selection was relaxed one generation. The base population was composed by 13 males and 83 females. The High (H) and Low (L) lines had 8 males and approximately 40 females each one per generation. Data from two generations of selection were used in this study.

Two rabbits (male and female) of the first parity of each doe were slaughtered at 9 weeks of age. Carcasses were prepared as recommended by the World Rabbit Science Association (Blasco and Ouhayoun, 1996). The following traits were recorded according to the methodology described by Hernández et al. (2004): live weight (LW), hot carcass weight (HCW), chilled carcass weight (CCW), reference carcass weight (RCW), scapular fat weight (SFW), perirenal fat weight (PFW), meat/bone ratio of hind leg (M/B), pH of LM, colour (lightness, L*; redness, a*; yellowness, b*) of the carcass and LM muscle. *Longissimus* muscles were excised from the carcass. Meat obtained from LM was ground, freeze-dried and scanned with a monochromator (model 5000, NIR Systems INC.) applying calibration equations previously developed (Zomeño et al., 2012).

Selection was made on the second parity using the average phenotypic IMF value of two full sibs of the first parity. Selection pressure on does was 18% for the first generation and 34% for the second generation. Males were selected from litters of sibs with the highest or lowest IMF value for H line and L line, respectively. Males were selected within sire families in order to reduce inbreeding.

Statistical Analysis

Data were analyzed using the following model:

$$y_{ijkl} = LG_i + SE_j + S_k + e_{ijkl}$$

where LG_i is the effect of line-generation (5 levels: base population, high line-generation 1, high line-generation 2, low line-generation 1 and low line-generation 2), SE_j is the effect of season (two levels: winter and summer), S_k is the effect of sex, and e_{ijkl} is the residual of the model.

A Bayesian analysis was performed. Bounded flat priors were used for all unknown parameters. Marginal posterior distributions for all unknown were estimated using Gibbs sampling. We used

a chain of 60,000 samples with a burn-in period of 10,000 for each trait. The following parameters were obtained from the marginal posterior distributions of the differences between lines: the median of the marginal posterior distribution of the difference (D); the highest posterior density region at 95% (HPD_{95%}); the probability of lines being different (probability of the difference between lines being higher than zero when this difference is higher than zero, or the probability of the difference between lines being lower than zero when this difference is lower than zero) (P); the lowest possible value of a difference with a probability of 95% (limit of the interval $[k, +\infty)$ when the difference is higher than zero) or the highest possible value with a probability of 95% (limit of the interval $(-\infty, k]$ when the difference is lower than zero) (k). Response to selection was estimated by the differences between H and L lines.

RESULTS AND DISCUSSION

Descriptive statistics of the traits are given in Table 1. The different adipose tissues represented a very small percentage of the chilled carcass (0.44 % for SF, 1.05 % for PF and 0.12% for IMF), which is in line with our previous results (Zomeño et al., 2010). LM showed a low lipid content (1.18 g/100g muscle) as expected, since it is one of the leanest muscles of the carcass (Pla et al., 2004). Values for carcass and meat colour variables and muscle pH are similar to those obtained in previous works (Hernández et al., 2004).

Table 1: Descriptive statistics of the traits analyzed.

Trait	N	Mean	SD	CVx100
LW (g)	502	1757	177	10.1
HCW (g)	502	1057	117	11.0
CCW (g)	502	1009	117	11.6
RCW (g)	502	797	97	12.1
SFW (g)	499	4.39	1.34	30.5
PFW (g)	502	10.6	4.52	42.6
IMF (g/100 g)	494	1.18	0.16	13.9
CL*	502	53.8	2.5	4.7
Ca*	502	3.13	0.84	26.8
Cb*	501	0.97	1.32	137
M/B	352	4.72	0.60	12.6
LML*	501	52.6	2.47	4.7
LMa*	501	3.66	0.99	26.9
LMb*	501	1.52	0.79	51.9
LMpH	497	5.53	0.10	1.8

N, number of animals; SD, standard deviation; CV, coefficient of variation; LW, live weight; HCW, hot carcass weight; CCW, chilled carcass weight; RCW, reference carcass weight; SFW, scapular fat weight; PFW, perirenal fat weight; IMF, intramuscular fat content of *Longissimus* muscle; CL*, Lightness of carcass surface; Ca*, redness of carcass surface; Cb*, yellowness of carcass surface; M/B, meat/bone ratio of hind leg; LML*, Lightness of *Longissimus* muscle; LMa*, redness of *Longissimus* muscle; LMb*, yellowness of *Longissimus* muscle; LMpH, ultimate pH of *Longissimus* muscle.

Table 2 shows the features of the estimated marginal posterior distributions of the differences between H and L lines for all traits studied. Intramuscular fat content was higher in line H than in line L ($P=1.00$), which represents a direct response to selection of 9.8%, of which 6.8% was obtained in the first generation and 3% in the second one. Selection experiments to increase IMF content have been previously conducted in pigs (Suzuki et al., 2005b; Schwab et al., 2009) and cattle (Sapp et al., 2002), with success.

Table 2: Features of the estimated marginal posterior distributions of the differences between High and Low lines (H-L) for intramuscular fat content (IMF) (g/100g), live weight, carcass and adipose tissues weights (g), meat to bone ratio of hind leg and muscle pH.

Trait	D	HPD _{95%}		P	k
IMF	0.11	0.06	0.15	1.00	0.07
LW	-64.6	-112	-16.7	1.00	-24.6
HCW	-33.9	-63.7	-1.92	0.98	-7.70
CCW	-32.6	-62.2	0.08	0.98	-5.62
RCW	-29.2	-56.0	-3.80	0.99	-8.03
SFW	-0.33	-0.71	0.05	0.95	0.00
PFW	-0.85	-2.02	0.53	0.91	0.20
C L*	-0.59	-1.37	0.10	0.95	0.00
C a*	0.13	-0.11	0.38	0.84	-0.07
C b*	0.06	-0.34	0.45	0.61	-0.28
M/B	-0.01	-0.18	0.18	0.55	0.14
LM L*	-0.42	-1.11	0.32	0.88	0.17
LM a*	0.19	-0.11	0.49	0.90	-0.06
LM b*	0.00	-0.24	0.22	0.50	-0.20
LM pH	-0.01	-0.03	0.03	0.61	0.02

D, median of the marginal posterior distribution of the difference H-L; HPD_{95%}, highest posterior density region at 95%; P , probability of the difference being greater than zero when $D>0$ and probability of the difference being lower than zero when $D<0$; k , limit of the interval [k , $+\infty$) when $D>0$ and $(-\infty, k]$ when $D<0$.

Live and carcass weights were slightly higher in line L than in line H ($P>0.98$), at least 24.6 g for LW and 5.62 g for CCW with a probability of 95% ($k=-24.6$ and -5.62 , respectively, table 2). Scapular and perirenal fat weights seem to be higher in line L ($P=0.95$ and $P=0.91$, respectively). Hence, there is a negative correlated response in LW and carcass weights but the evidence of correlated response in adipose tissue weights is rather weak. These results differed from those of Schwab et al. (2009) who obtained no correlated responses in growth performance traits and higher values of backfat measurements (mm) in a line selected for increased IMF. Genetic correlations between IMF and backfat thickness have been studied in pigs showing a wide range from 0.04 (reviewed by Sellier, 1998) to 0.64 (Solanes et al., 2009). The magnitudes of these correlations suggest that part of the genetic variation of IMF is independent of the genetic variation in overall lipid content of the carcass. In fact, Sapp et al. (2002) obtained no correlated response on fat thickness in bulls selected for IMF content suggesting that selection for increased IMF can be achieved without increasing carcass fat. Moreover, no phenotypic relationship

between IMF and perirenal fat was observed in a previous work in rabbits (Zomeño and Hernández, 2012). Maintenance energy is proportionally higher in rabbits than in pigs and cows due to its size; moreover rabbit is a particularly lean animal, and it is slaughtered at a rather early stage in comparison with beef and pigs. Thus it may happen that selection for IMF has no response on fat of depots, and it is possible that the positive correlation observed in other species only appears at later maturity states, in which fat content is higher. This means that genetics of IMF could be analyzed independently of the genetics of fat depots.

Line L had higher values for carcass L* ($P=0.95$) and lower values for carcass a*, although the probability of the differences for a* was lower ($P=0.84$) and the HPD_{95%} was larger (HPD_{95%} = [-0.11, 0.38]). No differences were observed between H and L lines for carcass b*. Line L also had higher values for muscle L* and lower for muscle a*, as it was observed in carcass. No differences were found between H and L lines for muscle b*, meat/bone ratio of hind leg and muscle pH. Similarly, Schwab et al. (2009) found that meat quality characteristics were almost not affected by selection for IMF.

CONCLUSIONS

Preliminary results show a response on IMF content after two generations of divergent selection for IMF, with a difference between High and Low lines of 0.11 g/100g. It seems to be a negative correlated response on live weight, carcass and adipose tissues weights. However, the evidence for adipose tissue is rather weak. Carcass and meat colour measurements as well as meat/bone ratio and muscle pH were almost not affected by selection.

ACKNOWLEDGEMENTS

This work was supported by AGL2008-05514-C02-01 project from the Spanish National Research Plan and PROMETEO/2009/125 project from the Generalitat Valenciana. C. Zomeño acknowledges a FPU grant from the MEC of Spain. Authors are grateful to V. Juste and F. Pardo for their technical assistance.

REFERENCES

- Blasco A., Ouhayoun J. 1996. Harmonization of criteria and terminology in rabbit meat research. Revised proposal. *World Rabbit Sci.*, 4, 93-99.
- Hernández P., Aliaga S., Pla M., Blasco A. 2004. The effect of selection for growth rate and slaughter age on carcass composition and meat quality traits in rabbits. *J. Anim. Sci.*, 82, 3138-3143.
- Laborda P., Mocé M. L., Santacreu M. A., Blasco A. 2011. Selection for ovulation rate in rabbits: Genetic parameters, direct response, and correlated response on litter size. *J. Anim. Sci.*, 89, 2981, 2987.
- Pla, M., Pascual, M., Ariño, B. 2004. Protein, fat and moisture content of retail cuts of rabbit meat evaluated with the NIRS methodology. *World Rabbit Sci.*, 12, 149-158.

- Saap R. L., Bertrand J. K., Pringle T. D., Wilson D. E. 2002. Effect of selection for ultrasound intramuscular fat percentage in Angus bulls on carcass traits of progeny. *J. Anim. Sci.*, 80, 2017, 2022.
- Schwab C. R., Baas T. J., Stalder K. J., Nettleton D. 2009. Results from six generation of selection for intramuscular fat in Duroc swine using real-time ultrasound. I. Direct and correlated phenotypic responses to selection. *J. Anim. Sci.*, 87, 2774-2780.
- Sellier P. 1998. Genetics of meat and carcass traits. In: Rothschild M.F., Ruvinsky A. (ed). *The Genetics of the Pig*. CAB Int., New York, USA, 463-510.
- Solanes, F.X., J. Reixach, M. Tor, J. Tibau, and J. Estany. 2009. Genetic correlations and expected response for intramuscular fat content in a Duroc pig line. *Liv. Sci.* 123, 63-69.
- Suzuki K., Irie M., Kadowaki H., Shibata T, Kumagai M., Nishida A. 2005a. Genetic parameter estimates of meat quality traits in Duroc pigs selected for average daily gain, longissimus muscle area, backfat thickness, and intramuscular fat content. *J. Anim. Sci.*, 83, 2058-2065.
- Suzuki K., Kadowaki H., Shibata T., Uchida H., Nishida A. 2005b. Selection for daily gain, loin-eye area, backfat thickness and intramuscular fat based on desired gains over seven generations of Duroc pigs. *Livest. Prod. Sci.*, 97, 193-202.
- Wood J.D., Enser M., Fisher A.V., Nute G.R., Sheard P.R., Richardson R.I., Hughes S.I., Whittington F.M. 2008. Fat deposition, fatty acid composition and meat quality: A review. *Meat Sci.*, 78, 343-358.
- Zomeño C., Blasco A., Hernández P. 2010. Influence of genetic line on lipid metabolism traits of rabbit muscle. *J. Anim. Sci.*, 88, 3419-3427.
- Zomeño C., Juste V., Hernández P. 2012. Application of NIRS for predicting fatty acids of intramuscular fat of rabbit. *Meat Sci. In Press*. DOI: 10.1016/j.meatsci.2012.01.009.
- Zomeño C., Hernández P. 2012. Relationships between lipid metabolism traits in carcass and muscle adipose tissues. *10th World Rabbit Congress. 2012, September, Sharm El-Sheikh, Egypt.*